

Xelstrym®
(dextroamphetamine) **II**
transdermal system

4.5mg/9 hours 9mg/9 hours 13.5mg/9 hours 18mg/9 hours

**XELSTRYM®: wearable
ADHD treatment that
lets you control**

Your time, Your way

*XELSTRYM is a once-daily patch that offers
flexibility to fit your changing schedule*

Work with your healthcare provider to determine
when to apply and when to remove XELSTRYM.
XELSTRYM should be removed within 9 hours.



**THE MINDFUL PATCH™ APP MAY HELP
YOU START AND STAY ON TRACK
WITH YOUR TREATMENT**



WHAT IS XELSTRYM?

XELSTRYM is a central nervous system stimulant prescription medicine used for the treatment of Attention-Deficit Hyperactivity Disorder (ADHD) in adults and children 6 years and older. It is not known if XELSTRYM is safe and effective in children younger than 6 years of age.

IMPORTANT SAFETY INFORMATION

Abuse, misuse, and addiction: XELSTRYM is a federally controlled substance (CII) because it contains dextroamphetamine that can be a target for people who abuse prescription medicines or street drugs. Keep XELSTRYM in a safe place to protect it from theft. Never sell or give your XELSTRYM to anyone else because it may cause death or harm to them and it is against the law. Tell your healthcare provider if you or your child have ever abused or been dependent on alcohol, prescription medicines, or street drugs.

Please read Important Safety Information throughout, [full Prescribing Information](#), including the BOXED WARNING, and the [Medication Guide](#).

What is XELSTRYM®?



The first and only amphetamine patch for ADHD

- XELSTRYM uses patch technology to deliver medication through the skin*—a method called transdermal drug delivery
- XELSTRYM has the same active metabolite (dextroamphetamine) as Vyvanse®†, a medication that has been used to treat ADHD since 2007



Treatment for ADHD should meet the demands of your changing schedule and offer flexibility

- XELSTRYM patch technology allows you and your healthcare provider to tailor your ADHD treatment to meet your daily needs
- Apply XELSTRYM 2 hours before the effect is needed and remove within 9 hours after application. Do not wear the patch for more than 9 hours
- Use only one patch per day (per 24 hours) and change application site when a new patch is applied

*Please see Instructions for Use at xelstry.com.

†There are no head-to-head safety or efficacy studies comparing Vyvanse® and XELSTRYM. VYVANSE® is a registered trademark of Takeda Pharmaceuticals U.S.A., Inc.

Xelstrym®
(dextroamphetamine) 
transdermal system
4.5mg/9 hours 9mg/9 hours 13.5mg/9 hours 18mg/9 hours



Patch not actual size.

IMPORTANT SAFETY INFORMATION (continued)

Do not use XELSTRYM if you or your child are:

- Allergic to amphetamine or any of the ingredients in XELSTRYM. See the end of the Medication Guide for a complete list of ingredients in XELSTRYM.

Please read Important Safety Information throughout, [full Prescribing Information](#), including the **BOXED WARNING**, and the [Medication Guide](#).

A treatment you can wear

The XELSTRYM[®] patch is clear,
discreet, and can be worn under your clothing
in one of 10 different application sites.



The Mindful Patch app offers customizable
application and removal reminders to help you
put your patch in the right spot and for the right
amount of time each day.

Upper back

Chest

Upper arm

Flank

Hip

Choose from right
or left side of body.

Four dosage strengths provide flexible options

Patches, US nickel,
and golf ball shown
at actual size.

18 mg/9 hrs
1.7 x 1.7 in

13.5 mg/9 hrs
1.5 x 1.5 in

9 mg/9 hrs
1.2 x 1.2 in

4.5 mg/9 hrs
.9 x .9 in

Golf ball = 1.7 in

Nickel = .8 in



Please read Important Safety Information throughout, [full Prescribing Information](#), including the BOXED WARNING, and the [Medication Guide](#).

Mindful Patch: A digital app for your ADHD treatment journey

Xelstrym®
(dextroamphetamine) 
transdermal system
4.5mg/9 hours 9mg/9 hours 13.5mg/9 hours 18mg/9 hours

The first and only app that helps you start and stay on track with your XELSTRYM® treatment over time. With the Mindful Patch app, you can:



Set and monitor your medication schedule

Customize your patch application and rotation schedules to help you stay on track with your treatment as determined by your healthcare provider.



Stay informed about your ADHD and wellbeing

With tracking of factors related to your ADHD and overall wellbeing, the Mindful Patch app gives you select insights about your ADHD journey that you can share with your healthcare provider.



Make the most of your treatment journey

Copay savings opportunities, including information about the Noven C.A.N.™ savings program.



Explore the Mindful Patch app



Download the
Mindful Patch app

Download on the
 App Store
Download via the
Apple store

GET IT ON
 Google Play
Download via the
Google app store

Please read Important Safety Information throughout, [full Prescribing Information](#), including the BOXED WARNING, and the [Medication Guide](#).

Savings and support are always within reach

Xelstrym®
(dextroamphetamine) 
transdermal system
4.5mg/9 hours 9mg/9 hours 13.5mg/9 hours 18mg/9 hours

Pay As Little As
\$25* at a Partner
Pharmacy

Xelstrym®
(dextroamphetamine) 
transdermal system
4.5mg/9 hours 9mg/9 hours 13.5mg/9 hours 18mg/9 hours

*Copay savings offer is available for commercially insured patients. At Noven C.A.N.™ Partner Pharmacies, eligible patients pay as little as \$25 for each XELSTRYM prescription. At non-Partner Pharmacies, eligible patients pay as little as \$50 for each XELSTRYM prescription. Patient is responsible for any remaining balance due after copay savings offer is applied. An offer is available for cash-paying patients. For eligibility information and full terms and conditions, visit xelstrym.com.

Care Access Network
NOVEN CAN

Get affordable access through the Copay Savings Program and Noven C.A.N.™ Partner Pharmacies.

Enrolling is simple! Sign up directly in the Mindful Patch app by texting “SAVE” to 515-517-6662, or by visiting XELSTRYM.com

IMPORTANT SAFETY INFORMATION (continued)

Do not use XELSTRYM if you or your child are:

- Allergic to amphetamine or any of the ingredients in XELSTRYM. See the end of the Medication Guide for a complete list of ingredients in XELSTRYM.
- Taking or have taken within the past 14 days a medicine called a monoamine oxidase inhibitor (MAOI), including the antibiotic linezolid or the intravenous medicine called methylene blue.

XELSTRYM can cause serious side effects, including:

- **Risks for people with serious heart disease.** Sudden death has happened in people who have heart defects or other serious heart disease. Your healthcare provider (HCP) should check you or your child carefully for heart problems before starting treatment with XELSTRYM. Tell your HCP if you or your child have any heart problems, heart disease, or heart defects. **Call your HCP or go to the nearest hospital emergency room right away if you or your child have any signs of heart problems such as chest pain, shortness of breath, or fainting during treatment with XELSTRYM.**

Please read Important Safety Information throughout, [full Prescribing Information](#), including the BOXED WARNING, and the [Medication Guide](#).

IMPORTANT SAFETY INFORMATION (continued)

XELSTRYM can cause serious side effects, including (continued):

- **Increased blood pressure and heart rate.** Your HCP should check you or your child's blood pressure and heart rate regularly during treatment with XELSTRYM.
- **Mental (psychiatric) problems.** Including new or worse behavior and thought problems; new or worse bipolar illness; new psychotic symptoms (such as hearing voices, or seeing or believing things that are not real); or new manic symptoms. Tell your HCP about any mental problems you or your child have or about a family history of suicide, bipolar illness, or depression. **Call your HCP right away if you or your child have any new or worsening mental symptoms or problems during treatment with XELSTRYM, especially hearing, seeing or believing things that are not real, or new manic symptoms.**
- **Slowing of growth (height or weight) in children.** Children should have their height and weight checked often while on XELSTRYM. Your HCP may stop treatment with XELSTRYM if your child is not growing or gaining height or weight as expected.
- **Circulation problems in fingers and toes (peripheral vasculopathy, including Raynaud's phenomenon).** Fingers or toes may feel numb, cool, or painful. Fingers or toes may change color from pale, to blue, to red. Tell your HCP if you or your child have any numbness, pain, skin color change, or sensitivity to temperature in the fingers or toes. **Call your HCP right away if you or your child have any signs of unexplained wounds appearing on fingers or toes while using XELSTRYM.**
- **Serotonin syndrome.** Symptoms such as agitation, fast heartbeat, flushing, seizures, coma, sweating, loss of coordination, confusion, dizziness, tremors, stiff muscles, muscle twitching, seeing or hearing things that are not real (hallucinations), changes in blood pressure, high body temperature (hyperthermia), nausea, vomiting, or diarrhea may occur. Stop using XELSTRYM and call your HCP or go to the emergency room if symptoms occur. Serotonin syndrome may occur if XELSTRYM is taken with certain medicines and may be life-threatening.
- **Allergic skin rash (contact sensitization).** Stop using XELSTRYM and tell your HCP right away if you or your child develop swelling or blisters at or around the application site.
- **Application site reactions.** Reactions have happened while wearing XELSTRYM and after removing it. Symptoms include pain, itching, burning feeling, redness, discomfort, or swelling at the application site. Call your HCP if you or your child develop any application site reactions that do not resolve on their own.
- **New or worsening tics or worsening Tourette's syndrome.** Tell your HCP if you or your child get any new or worsening tics or worsening Tourette's syndrome during treatment with XELSTRYM.

Please read Important Safety Information throughout, [full Prescribing Information](#), including the BOXED WARNING, and the [Medication Guide](#).



IMPORTANT SAFETY INFORMATION (continued)

Before taking XELSTRYM tell your HCP about all the medications you take and any medical conditions including if you:

- Are pregnant or plan to become pregnant. It is not known if XELSTRYM may harm your unborn baby. There is a pregnancy registry for women who are exposed to XELSTRYM during pregnancy. The purpose of the registry is to collect information about the health of women exposed to XELSTRYM and their babies. For more information on registering with the National Pregnancy Registry for Psychiatric Medications call 1-866-961-2388 or visit online at <https://womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry/othermedications/>
- Are breastfeeding or plan to breastfeed. XELSTRYM can pass into your milk. Do not breastfeed while taking XELSTRYM. Talk to your doctor about the best way to feed your baby if you take XELSTRYM.

What should I avoid while using XELSTRYM?

- Do not drive, operate heavy machinery, or do other potentially dangerous activities until you know how XELSTRYM affects you.
- Avoid exposing the application site to direct external heat sources such as hair dryers, heating pads, electric blankets, heat lamps, saunas, hot tubs, and heated water beds, as exposure to heat can cause too much medicine to pass into your body and cause serious side effects.

The most common side effects of XELSTRYM include:

- | | | |
|----------------------|----------------------------|------------------------|
| • decreased appetite | • increased blood pressure | • vomiting |
| • trouble sleeping | • headache | • irritability |
| • stomach pain | • muscle twitching (tics) | • increased heart rate |
| • nausea | | |

Please read Important Safety Information throughout, [full Prescribing Information](#), including the BOXED WARNING, and the [Medication Guide](#).

You are encouraged to report negative side effects of prescription drugs to FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.



XELSTRYM® is a once-daily ADHD patch designed to fit into your daily schedule

The patch should not be worn for more than 9 hours.

- The Mindful Patch app may help you start and stay on track with XELSTRYM
- To learn more about XELSTRYM and to access a library of helpful resources—including the Instructions for Use—download the **Mindful Patch app** or visit XELSTRYM.com



Download the
Mindful Patch app



Download on the
App Store

Download via the
Apple store



GET IT ON
Google Play

Download via the
Google app store

Xelstrym®
(dextroamphetamine) 
transdermal system
4.5mg/9 hours 9mg/9 hours 13.5mg/9 hours 18mg/9 hours



IMPORTANT SAFETY INFORMATION (continued)

Do not use XELSTRYM if you or your child are:

- Allergic to amphetamine or any of the ingredients in XELSTRYM. See the end of the Medication Guide for a complete list of ingredients in XELSTRYM.
- Taking or have taken within the past 14 days a medicine called a monoamine oxidase inhibitor (MAOI), including the antibiotic linezolid or the intravenous medicine called methylene blue.

Please read Important Safety Information throughout, [full Prescribing Information](#), including the BOXED WARNING, and the [Medication Guide](#).



XELSTRYM® is a trademark of Noven Therapeutics, LLC.
Mindful Patch™ is a trademark of Noven Therapeutics, LLC.
Noven C.A.N. and its related logo are trademarks of Noven Therapeutics, LLC.
©2024 Noven Therapeutics, LLC. All rights reserved. For US audience only.
NPX-3002-16 12/2024